

General Practice Referral Form

Email: life@lifeprogram.org.au or **Fax:** (03) 96671757

| teferral Date | | Does your patient speak English? | Yes | | | |
|---|--|--|--|--|--|--|
| lame | | What is the main language spoken at home? | | | | |
| address | | Aboriginal or Torres Strait Islander descent? | | | | |
| | | Past medical history | | | | |
| itate | Postcode | | | | | |
| hone | | | | | | |
| mail | | | | | | |
| OOB | Gender | Current smoker? Yes No | | | | |
| Country of birth | | Blood pressure systolic/ | diastol | | | |
| Cultural background | | Waist circumference (cm) | | | | |
| | ria: • Diahetes • Active Cancer • C | Cardiovascular Disease (CVD) diagnosis in the last 3 r | months | | | |
| | ne suitability of the <i>Life!</i> program f | | | | | |
| | | | | | | |
| Patient eligibility | | | | | | |
| Choose <u>one</u> of the following | | | | | | |
| Α | В | С | | | | |
| ≥ 18 years and AUSDRISK ≥ 12 and BMI ≥ 25kg/m² | | ≥ 18 years with one or more of the following pre-existing conditions (Please tick 🗸 and document) | | | | |
| (if patient self-identifies | and/or Torres Strait | | | | | |
| as being of Asian background a BMI ≥ | Islander descent and have an intermediate | Cardiovascular Disease* Gestational Diabetes Chronic Kidney Disease Pre-diabetes (IFG or IGT) | | | | |
| 23kg/m² is accepted)* | or high risk score when | | | | | |
| AUSDRISK Score | referred by a GP clinic. | Polycystic Ovary Syndrome Familial Hyperch | olesterolemia | | | |
| Height (cm) | CVD risk | Serum total cholesterol > 7.5mmol/L (initial reading | g) | | | |
| Weight (kg) | category | Syst BP of ≥ 180 mmHg or Diast BP ≥110mmHg (read | ling) | | | |
| BMI | | | | | | |
| MBS items apply for a health | The following time-based MBS item applies for a heart health | or further information or clarification, please refer to the definitions pag | ge. | | | |
| assessment 701, 703, 705, 707, 715 | check: 699,177 | | | | | |
| | | | | | | |
| Deferrer details | | | | | | |
| Referrer details | | Completing the referral – $\sqrt{\text{tick to confirm}}$ | | | | |
| Referrer details | | Completing the referral − ✓ tick to confirm Yes, patient consents to be enrolled in the | program | | | |
| | | Yes, patient consents to be enrolled in the | | | | |
| lame | | , , | | | | |
| Jame Clinic | | Yes, patient consents to be enrolled in the Blood pathology report within 12 months o | | | | |
| lame Clinic Address | Postcode | Yes, patient consents to be enrolled in the Blood pathology report within 12 months of Fasting blood glucose (FBG) or HbAlc | f referral date* | | | |
| lame Clinic Address | Postcode | Yes, patient consents to be enrolled in the Blood pathology report within 12 months o Fasting blood glucose (FBG) or HbA1c Lipid Profile (TC/TRIG/HDL/LDL) | f referral date* | | | |
| Jame Clinic Address State Phone | Postcode | Yes, patient consents to be enrolled in the Blood pathology report within 12 months o Fasting blood glucose (FBG) or HbAlc Lipid Profile (TC/TRIG/HDL/LDL) Completed AUSDRISK form is using criteria | f referral date* | | | |
| Jame Clinic Address State Chone ax | Postcode | Yes, patient consents to be enrolled in the Blood pathology report within 12 months of Fasting blood glucose (FBG) or HbAlc Lipid Profile (TC/TRIG/HDL/LDL) Completed AUSDRISK form is using criterial Diabetes excluded* Yes, patient is happy for Life! to contact the or about their experience in the program. Yes, patient is happy for Life! to contact the second contact their experience in the program. | of referral date* A em for research em to discuss | | | |
| Jame Clinic Address State Phone | Postcode | Yes, patient consents to be enrolled in the Blood pathology report within 12 months of Fasting blood glucose (FBG) or HbA1c Lipid Profile (TC/TRIG/HDL/LDL) Completed AUSDRISK form is using criterial Diabetes excluded* Yes, patient is happy for Life! to contact the or about their experience in the program. | of referral date* A em for research em to discuss | | | |

A. AUSDRISK + BMI

The Australian type 2 diabetes risk assessment tool (AUSDRISK)

is a short list of questions to help assess a person's risk of developing type 2 diabetes over the next five years.

https://www.health.gov.au/resources/apps-and-tools/the-australian-type-2-diabetes-risk-assessment-tool-ausdrisk

Life! waist circumference/clothing size chart for males and females.

https://lifeprogram.org.au/wp-content/uploads/2021/07/Waist-Measurement-Flyer.pdf

Body mass index (BMI) is one way to assess whether a person's weight is in the healthy range. The BMI combines a person's height and weight to form a measure that can help predict their risk of developing disease.

Body mass index = $\frac{\text{weight (kg)}}{\text{height (m)}^2}$

www.aihw.gov.au/reports/overweight-obesity/overweight-and-obesity/contents/about

The following time-based MBS items apply for a health assessment: 701, 703, 705, 707, 715

Asian sub-regions

Asian background includes three distinct sub-regions (Parliament of Australia, 1997):

Northeast Asia (Mainland China, Hong Kong, Japan, Korea, Taiwan);

Southeast Asia (Brunei, Cambodia, Indonesia, Laos, Malaysia, Myanmar, the Philippines, Singapore, Thailand, Vietnam); and Southern Asia (Afghanistan, Bangladesh, India, Pakistan, Sri Lanka, Nepal).

B. Cardiovascular risk score

The Australian CVD risk calculator combines several risk factors to calculate a risk score (expressed as a percentage), which is a person's chance of having a CVD event such as a heart attack or stroke in the next five years.

www.cvdcheck.org.au

C. Pre-existing conditions

Cardiovascular disease*1

- Acute rheumatic fever/ rheumatic heart disease
- Angina
- Angioplasty
- Arrhythmias/Atrial fibrillation
- Cardiomyopathy
- · Chronic Heart Failure
- · Coronary Heart Disease
- · Myocardial infarction
- · Peripheral Vascular Disease
- Stroke

*Must not have been diagnosed within the last 3 months.

*Must be medically stable and must not have had a hospital admission in the last 12 months for the condition.

Gestational diabetes

To participate in the program, women must not be pregnant.

Moderate or severe kidney disease²

- Persistent proteinuria or GFR < 45mL/min/1.73m² or
- Stage 3b chronic kidney disease or greater

Pre-diabetes3456

Blood glucose levels are higher than normal but not high enough to be classified as diabetes.

Impaired Fasting Glucose (IFG)

Blood glucose levels are escalated in the fasting state but not high enough to be classified as diabetes. It is diagnosed on the basis of a result between 6.1 and 6.9 mmol/L

• Impaired Glucose Tolerance (IGT)

Blood glucose level is higher than normal two hours after an OGTT but not high enough to be diagnosed with diabetes. Two-hour plasma glucose is between 7.8 and 11.0 mmol/L. Fasting blood glucose level may still be in the normal range.

Glycated haemoglobin (HbAlc)
 Pre-diabetes can be diagnosed with a HbAlc of 6.0-6.4%

Familial hypercholesterolemia⁷

The Dutch Lipid Clinic Network Score Criteria required for diagnosis.

Serum cholesterol >7.5 mmol/L

If on medication managing the condition, patient is still eligible for the program.

Systolic BP ≥ 180mmHg or diastolic BP ≥ 110mmHg8

If on medication managing the condition, patient is still eligible for the program.

Polycystic Ovarian Syndrome⁹

Rotterdam diagnostic criteria. Requires two of:

- · Oligo- or anovulation
- · Clinical and/or biochemical hyperandrogenism
- Polycystic ovaries on ultrasound (and exclusion of other aetiologies)

Blood Test Results¹⁰

A patient's blood test result is required to ensure that type 2 diabetes has been excluded. Please attach patient blood test results within the last 12 months of the *Life!* referral date. If the patient is currently pregnant and has been diagnosed with GDM, post partum results will be requested once the patient is enrolled in the program.

- Full Blood Lipids
- Fasting Blood Glucose (FBG) or HbA1c

Further testing to exclude type 2 diabetes

If FBG is 5.5 - 6.9 mmol/L an Oral Glucose Tolerance Test (OGTT) is recommended.

These are via venous sampling under laboratory methodology.

Diagnostic criteria for type 2 diabetes (RACGP clinical guidelines 2016-2018)¹¹

Patient is excluded from the program if they meet one or more of the following:

- Fasting blood glucose (FBG) >7.0 mmol/L or random blood glucose >11.1 mmol/L confirmed by a second abnormal FBG on a separate day
- Oral glucose tolerance test (OGTT) before (fasting) and two hours after an oral 75 g glucose load is taken. Blood glucose is measured. Diabetes is diagnosed as FBG ≥7.0 mmol/L or twohour blood glucose ≥11.1 mmol/L
- Glycated haemoglobin (HbAlc) ≥48 mmol/mol (6.5%; on two separate occasions)

These are via venous sampling under laboratory methodology.

Additional information

If you refer your patient while they are pregnant, please note that they will not be able to commence the program until they are at least 6 weeks postpartum. Both the referrer and the patient will be updated on the progress of the referral.

- 1. Heart Foundation www.heartfoundation.org.au/bundles/your-heart
- 2. Kidney Health kidney.org.au/health-professionals/ckd-management-handbook
- 3. Diabetes Australia www.diabetesaustralia.com.au/about-diabetes/pre-diabetes/
- 4. National Diabetes Services Scheme www.ndss.com.au/understandina-pre-diabetes
- Royal Australian College of General Practitioners LTD (RACGP) www.racgp.org.au /clinical-resources/clinical-guidelines/key-racgp-guidelines/view-all-racgp-guidelines/ guidelines-for-preventive-activities-in-general-pr/preamble/introduction
- RACGP www.racgp.org.au/your-practice/guidelines/diabetes/3-screening,-riskassessment,-case-finding-and-diagnosis/33-impaired-fasting-glucose-or-impairedglucose-tolerance/
- Familial Hypercholesterolemia Australasia Network www.athero.org.au/fh/patients/ diagnosis/
- 8. **Heart Foundation** www.heartfoundation.org.au/bundles/for-professionals/hypertension
- 9. **Jean Hailes for Women's Health** www.jeanhailes.org.au/health-a-z/pcos
- RACGP https://www.racgp.org.au/clinical-resources/clinical-guidelines/key-racgpguidelines/view-all-racgp-guidelines/diabetes/preventing-progression-to-type-2-diabetes
- RACGP https://www.racgp.org.au/clinical-resources/clinical-guidelines/key-racgpguidelines/view-all-racgp-guidelines/diabetes/preventing-progression-to-type-2-diabetes





The Australian Type 2 Diabetes Risk Assessment Tool (AUSDRISK)

The AUSDRISK tool estimates your risk of getting Type 2 Diabetes in the next 5 years.

Not everyday

| 1. | Your age group: | | 9. On average, wou | ld you say you do at le | ast 2.5 hours of | | |
|----|---|----------|--|---|-----------------------|--|--|
| | Under 35 years 0 points | | physical activity | per week (for example | , 30 minutes a | | |
| | 35-44 years | 2 points | day on 5 or more days a week)? | | | | |
| | 45-54 years | 4 points | Yes | | 0 points | | |
| | 55-64 years | 6 points | No | | 2 points | | |
| | 65 years or over | 8 points | | surement taken below the ribs level of the navel, while standing): | | | |
| 2. | Your gender: | | Waist measurem | ent (cm) | | | |
| | Female | 0 points | 11. Complete either the purple or orange box below, as appropriate: | | | | |
| | Male | 3 points | | | | | |
| 3. | Your ethnicity/country of birth: | | For those of Asian or Aboriginal or Torres Strait | | | | |
| | . Are you of Aboriginal, Torres Strait Islander, Pacific Islander or Maori descent? | | Islander descent: Men | Women | | | |
| | No | 0 points | Less than 90cm | Less than 80cm | 0 points | | |
| | Yes | 2 points | 90-100cm | 80-90cm | 4 points | | |
| | | 2 points | More than 100cm | More than 90cm | 7 points | | |
| | Where were you born? | | | | | | |
| | Australia | 0 points | For all others: | | | | |
| | Southern Europe | 2 points | Men | Women | | | |
| | North Africa | 2 points | Less than 102cm | Less than 88cm | 0 points | | |
| | Asia (including the Indian sub-continent) | 2 points | 102-110cm | 88-100cm | 4 points | | |
| | Middle East | 2 points | More than 110cm | More than 100cm | 7 points | | |
| | Other | 0 points | | | | | |
| 4. | Have either of your parents, or any of your brothers or sisters been diagnosed with diabetes (type 1 or type 2)? Add up your points | | | | | | |
| | No | 0 points | Your risk of develop | ing type 2 diabetes wit | hin 5 years*: | | |
| | Yes | 3 points | E av lage. Lavy viels | | | | |
| 5. | Have you ever been found to have high blood glucose (sugar) for example, in a health examination, during an illness, during pregnancy? 5 or less: Low risk Approximately one person in every 100 will develop diabetes. | | | | | | |
| | No | 0 points | 6–11: Intermediate risk | | | | |
| | Yes | 6 points | For scores of 6–8, approximately one person in every | | | | |
| 6. | Are you currently taking medication for high blood pressure? | | 50 will develop diabetes. For scores of 9–11, approximately one person in every 30 will develop diabetes. | | | | |
| | No | 0 points | 12 or more: High risk | | | | |
| | Yes | 2 points | | | | | |
| 7. | Do you currently smoke cigarettes or any tobacco products on a daily basis? | other | 14 will develop dia | i, approximately one pe betes. I, approximately one pe | · | | |
| | No | 0 points | seven will develop diabetes. | | | | |
| | Yes | 2 points | For scores of 20 and above, approximately one person in every three will develop diabetes. | | | | |
| 8. | How often do you eat vegetables or fruit? | | , | · | ا د داد ما ما داد داد | | |
| | Everyday | 0 points | *The overall score m people younger that | ay overestimate the ris ın 25. | k of alabetes in | | |
| | | | 1 1 / | | | | |

__ 1 point

The Australian Type 2 Diabetes Risk Assessment Tool was originally developed by the International Diabetes Institute on behalf of the Australian, State and Territory Governments as part of the COAG Diabetes reducing the risk of type 2 diabetes initiative.